

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

JN




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification⁶ : C07K 7/00</p>	<p>A2</p>	<p>(11) International Publication Number: WO 98/34949 (43) International Publication Date: 13 August 1998 (13.08.98)</p>
<p>(21) International Application Number: PCT/EP98/00599 (22) International Filing Date: 4 February 1998 (04.02.98) (30) Priority Data: FI97A000020 7 February 1997 (07.02.97) IT (71) Applicant (for all designated States except US): MENARINI RICERCHE S.P.A. [IT/IT]; Via Tito Speri, 10, I-00040 Pomezia (IT). (72) Inventors; and (75) Inventors/Applicants (for US only): GIORGI, Raffaello [IT/IT]; Via Delle Piagge, 9, I-56124 Pisa (IT). DI BUGNO, Cristina [IT/IT]; Via R. Sanzio, 16, I-56122 Pisa (IT). GIANNOTTI, Danilo [IT/IT]; Via Roma, 128, I-55011 Altopascio (IT). MAGGI, Carlo, Alberto [IT/IT]; Via Michelazzi, 43, I-50141 Firenze (IT). (74) Agent: PASSINI, Angelo; Notarbartolo & Gervasi S.p.A., Corso di Porta Vittoria, 9, I-20122 Milan (IT).</p>		<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p>Published Without international search report and to be republished upon receipt of that report.</p>

(54) Title: MONOCYCLIC COMPOUNDS WITH FOUR BIFUNCTIONAL RESIDUES HAVING NK-2 ANTAGONIST ACTION

(57) Abstract

The present invention refers to compounds of general formula (I) having NK-2 antagonist action, pharmaceutical compositions containing them, and processes for their preparation.



FOR THE PURPOSES OF INFORMATION ONLY


Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

MONOCYCLIC COMPOUNDS WITH FOUR BIFUNCTIONAL RESIDUES HAVING NK-2 ANTAGONIST ACTION

Scope of the invention

The present invention refers to new compounds having the general formula (I):



in which:

X_1, X_2, X_3, X_4 , which may be the same or different from one another, represent a group chosen from among $-\text{CONR}-$, $-\text{NRCO}-$, $-\text{OCO}-$, $-\text{COO}-$, $-\text{CH}_2\text{NR}-$, $-\text{NR}-\text{CH}_2-$, CH_2-CH_2 , where R is H or a C_{1-3} alkyl or benzyl;

f, g, h, m , which may be the same or different from one another, represent a number chosen from among 0, 1 or 2;

R_1 and R_2 , which may be the same or different from one another, represent a $-(\text{CH}_2)_r-\text{Ar}$ group, where $r = 0, 1, 2$ and where Ar is an aromatic group chosen from among: benzene, naphthalene, thiophene, benzothiophene, pyridine, quinoline, indole, furan, benzofuran, thiazole, benzothiazole, imidazole, and benzo-imidazole, the said Ar group being possibly substituted with a maximum of 2 residues chosen from among C_{1-3} alkyl or halo-alkyl, C_{1-3} alkoxyl, C_{2-4} amino-alkoxyl, halogen, OH, NH_2 , $\text{NR}_{13}\text{R}_{14}$ where R_{13} and R_{14} , which may be the same or different from one another, represent hydrogen or C_{1-3} alkyl;

R_3 represents a group chosen from among:

- hydrogen

- linear or branched alkyl having the formula $\text{C}_n\text{H}_{2n+1}$, with $n = 1-5$, cyclo-alkyl or alkylcyclo-alkyl groups having the formula $\text{C}_n\text{H}_{2n-1}$ with $n = 5-9$

- $(\text{CH}_2)_r-\text{Ar}_1$, where $r = 0, 1, 2$ and where Ar_1 is an aromatic group chosen from among: benzene, naphthalene, thiophene, benzothiophene, pyridine, quinoline, indole, furan, benzofuran, thiazole, benzothiazole, imidazole, and benzo-imidazole, the said Ar_1 group being possibly substituted with a maximum of 2

residues chosen from among C_{1-3} alkyl or halo-alkyl, C_{1-3} alkoxy or amino-alkoxy, halogen, OH, NH_2 , $NR_{13}R_{14}$, where R_{13} and R_{14} , which may be the same or different from one another, represent hydrogen or C_{1-3} alkyl;

R_4 represents a group chosen from among:

5 - hydrogen or C_{1-6} alkyl

- L-Q, where L is a chemical bond or a linear or branched C_{1-6} alkyl residue and Q is a group chosen from among:

i) H, OH, OR_9 , NH_2 , NR_9R_{10} , guanidine, sulphate, phosphonate, phosphate, where R_9 and R_{10} , which may be the same or different from one another, represent a hydrogen, C_{1-3} alkyl group, C_{1-3} hydroxyalkyl, C_{1-3} dihydroxyalkyl, C_{1-3} alkyl-CONHR₁₂, C_{1-3} alkyltetrazole, C_{1-3} alkyl-COOH or wherein R_9R_{10} joined together form with the N-atom a saturated 4-6 membered heterocycle possibly containing a further heteroatom chosen in the group consisting of N, O, S and wherein R_{12} is a mono-, di-, tri-glycosidic group possibly protected with one or more C_{1-3} -acyl groups or substituted with amino-groups or C_{1-3} acylamino-groups;

ii) COOH, tetrazole, SO_2NH_2 , $SO_2NHCOOR_8$, CONHR₈, NHCOR₈, where R_8 represents a linear or cyclic C_{1-6} alkyl chain containing one or more polar groups chosen from among the group: OH, NH_2 , $NR_{15}R_{16}$, COOH, CONHR₁₂, PO_3H , SO_3H , OR_{11} and where R_{15} and R_{16} , which may be the same or different from one another, represent a hydrogen or C_{1-3} alkyl group, and where R_{11} is a C_{1-3} alkyl or C_{2-4} amino-alkyl chain, R_{12} is a mono-, di-, tri-glycosidic group possibly protected with one or more C_{1-3} acyl groups or substituted with amino-groups or C_{1-3} acylamino-groups or $R_{15}R_{16}$ joined together form with the N-atom a saturated 4-6 membered heterocycle possibly substituted with C_{1-3} alkyl-groups or with saturated 4-6 membered heterocycle-groups containing at least an N-atom;

iii) COOR₁₇, CONHR₁₂, OR₁₂ where R_{12} is a mono-, di- or tri-glycoside group possibly protected with one or more C_{1-3} acyl groups or substituted with amine or C_{1-3} acylamine groups and R_{17} is a group R_{12} as above defined or a group C_{1-3} alkyl, C_{1-3} alkylphenyl, wherein the phenyl-group can be substituted with a group OH, NO_2 , NH_2 , CN, CH_3 , Cl, Br;

R_5 , R_6 , R_7 , which may be the same or different from one another, represent a hydrogen or C_{1-3} alkyl group.

Also included in the present invention are the pharmaceutically acceptable salts, the processes for their preparation, and the pharmaceutical compositions
5 containing them.

In view of the presence of chiral centres in the compounds of formula (I), also the individual enantiomers and their mixtures, both in the racemic form and in the non-racemic form, form part of the present invention.

State of the art

10 The NK-2 receptor of tachykinins is widely expressed in the peripheral nervous system of mammals. One of the various effects produced by the selective stimulation of the NK-2 receptor is the contraction of smooth muscle. Hence antagonists of the NK-2 receptor may be considered agents capable of controlling excessive contraction of smooth muscle in any pathological
15 condition in which the release of tachykinins concurs in the genesis of the corresponding disorder.

In particular, the bronchospastic and inflammatory component of asthma, coughing, pulmonary irritation, intestinal spasms, spasms of the biliary tract, local spasms of the bladder and of the ureter during cystitis, kidney infections

It has now surprisingly been found that products of lower molecular weight, monocyclic ones, containing only four bifunctional residues linked together via peptide or pseudopeptide bond, present high pharmacological activity associated to a considerable selectivity for the human NK-2 receptor, and thus are proposed as valid alternatives.

Detailed description of the invention

The present invention therefore sets itself the aim of making available new monocyclic compounds containing four bifunctional residues linked together via peptide or pseudopeptide bonds having antagonistic action on the NK-2 receptor, with the general formula (I), as defined previously.

Also forming part of the present invention are the pharmaceutically acceptable salts, the processes for their preparation, and the pharmaceutical compositions containing them.

In view of the presence of chiral centres in the compounds of formula (I), also the individual enantiomers and their mixtures, both in the racemic form and in the non-racemic form, form part of the present invention.

According to the invention preferred compounds of general formula (I) are those in which:

f, g, h, m, which may be the same or different from one another, may be 0 or 1;
R₁ and R₂, which may be the same or different from one another, represent the side chain of a natural amino acid chosen from among tryptophan, phenyl alanine, tyrosine, histidine or the side chain of a non-natural amino acid chosen in the group:

tryptophan and phenyl alanine, either mono- or di-substituted with residues chosen from among C₁₋₃ alkyl or halo-alkyl, C₁₋₃ alkoxy or amino-alkoxy, halogen, OH, NH₂, NR₁₃R₁₄, where R₁₃ and R₁₄, which may be the same or different from one another, represent a hydrogen or C₁₋₃ alkyl group;

R₃ represents a group chosen from among:

- linear or branched alkyl having the formula C_nH_{2n+1}, with n = 1-5 (chosen in the group consisting of methyl, ethyl, propyl, isopropyl, n-butyl, t-butyl) cycloalkyl or alkylcycloalkyl of formula C_nH_{2n-1} with n = 5-9 (chosen in the group consisting of cyclopentyl, cyclohexyl, methylcyclohexyl)

- $(\text{CH}_2)_r\text{-Ar}_1$, where $r = 1$ or 2 and where Ar_1 is an aromatic group chosen from among: α -naphthyl, β -naphthyl, phenyl, indole, the said Ar_1 group being possibly substituted with a maximum of 2 residues chosen from among C_{1-3} alkyl, CF_3 , C_{1-3} alkoxy, Cl, F, OH, NH_2 ;

5 R_4 represents an L-Q group where:

L is a chemical bond or CH_2 , and

Q is a group chosen from among:

- OH, NH_2 , NR_9R_{10} , OR_{11} , and where R_9 and R_{10} , which may be the same or different from one another, represent a hydrogen or C_{1-3} alkyl group, C_{1-3} hydroxy alkyl, C_{1-3} dihydroxyalkyl, C_{1-3} alkyl- CONHR_{12} wherein R_{12} is a monoglycosidic group derived from D or L pentoses or hexoses (chosen in the group consisting of ribose, arabinose, glucose, galactose, fructose, glucosamine, galactosamine and their N-acetylated derivatives), C_{1-3} alkyltetrazole, C_{1-3} alkyl-COOH or wherein R_9R_{10} are joined together to form with the N atom a morpholine or a
- 10 piperidine ring and where R_{11} is a C_{1-3} alkyl chain, or a C_{2-4} amino-alkyl chain;
- NHCOR_8 wherein R_8 is a cyclohexane containing from 2 to 4 OH groups, a C_{1-6} alkylchain containing a polar group (chosen in the group consisting of NH_2 , COOH, CONHR_{12} (wherein R_{12} is as hereabove defined) or [1,4']bipiperidine)
- COOH, COOR_{17} or CONHR_{12} , wherein R_{12} is as hereabove defined and R_{17} is
- 20 as R_{12} or a group 4-nitrobenzyl and R_{12} is a monoglycosidic group derived from D or L pentoses or hexoses (chosen in the group consisting of ribose, arabinose, glucose, galactose, fructose, glucosamine, galactosamine and their N-acetylated derivatives).

- R_5 , R_6 , R_7 are H.

25 Likewise preferred are isomers that present an R configuration on the carbon atom that carries the R_3 and R_7 substituents.

Pharmaceutically acceptable salts of compounds of formula (I) include the salts with inorganic acids (such as, hydrochloric acid, hydrobromic acid, hydrogen iodide, sulphuric acid, nitric acid, phosphoric acid) and organic acids (such as, acetic, propionic, succinic, malonic, citric, tartaric, metasulphonic, *para*-toluenesulphonic acids), as well as salts of pharmaceutically acceptable bases,

30 both inorganic (such as, hydroxides of sodium, potassium, calcium,

magnesium, zinc and aluminium) and organic bases (such as, amines like methyl amine, diethyl amine, triethyl amine, ethyl amine, tromethamine or piperidine).

According to the invention, the compounds of formula (I) containing peptide or
5 pseudopeptide bonds may be obtained by means of classical condensation methods using techniques known in the literature. The general method chosen by us for preparing the peptide compounds ($X_1-X_4 = -CONR-$, $-NRCO-$) involves the synthesis in solution of the linear peptide chain using amino acids, dicarboxyl or diamine derivatives suitably protected and, after selective
10 deprotection of the C- and N-terminals, cyclization in polar organic solvents in diluted solution. As method of activation of the carboxyl groups, that using PyBOP and DIEA in DMF or HBT and EDC in DMF are generally preferred.

To provide an example, the attached diagram presents the general synthesis of compounds of formula (I) in which $X_1 = X_2 = X_3 = X_4 = -CONH-$.

15 The dicarboxyl precursors 7 containing the R_4 group, and the diamine precursors 4 containing the R_3 and R_7 groups were prepared using methods described in the literature.

In particular, the synthesis of the succinic derivatives, with $R_4 = \text{alkyl or } (CH_2)_n-\text{Ar}$, is described by R. Conrow *et al.*, J. Org. Chem., 1986, 51, 938 and by S.G.
20 Cohen *et al.*, J. Am. Chem. Soc., 1968, 90, 3495, whilst in the case of $R_4 = \text{H}$, amine group, hydroxyl group or carboxyl group, the following were respectively used: succinic anhydride, aspartic acid, malic acid or carboxysuccinic acid appropriately protected.

The synthesis of the ethylene diamine derivatives containing the R_3 , R_7 groups
25 was performed starting from the corresponding N-protected amino acids by reduction of the carboxyl to alcohol with $BH_3.THF$ (C.F. Stanfield *et al.*, J. Org. Chem., 1981, 46, 4797, 4799; I.R. Ollmann *et al.*, Bioorg. Med. Chem., 1995, 3, 969), conversion to azide via the corresponding mesylate and subsequent reduction to amino group (P.G. Mattingly, Synthesis, 1990, 366; P.M. O'Brien *et al.*, J. Med. Chem., 1994, 37, 1810).
30

The compounds containing reduced peptide bonds ($X_1-X_4 = -CH_2-NR-$, $-NR-CH_2-$) were synthesized in solution according to known methods, such as

reductive amination of the aldehyde of an amino acid with the amine function of a protected amino acid or peptide, in the presence of NaCNBH_4 as reducer in DMF/AcOH (K.A. Jacobson *et al.*, J. Med. Chem., 1983, 26, 492; R.F. Borch *et al.*, J. Am. Chem. Soc., 1971, 93, 2897; J.P. Salvi *et al.*, Tetr. Lett., 1994, 35, 1181). The aldehydes were obtained by reduction with LiAlH_4 of the
5 corresponding protected amino acids, N,O-dimethylhydroxy-amates according to the method described by J.A. Feherentz *et al.*, Synthesis, 1983, 676 and Int. J. Peptide Res., 1985, 26, 236.

The compounds of formula 10 wherein R_4 is NH_2 or COOH can be derivatized
10 into compounds of formula 1 wherein R_4 is NR_9R_{10} , guanidine, tetrazole, NHCOR_8 , CONHR_8 , COOR_{17} , CONHR_{12} , wherein R_9 , R_{10} , R_8 , R_{12} and R_{17} are as above defined, according to known methods.

The compounds of formula (I) as specified above have proved to be powerful antagonists of the NK-2 receptor of tachykinins, and hence can be administered
15 as agents capable of controlling any central or peripheral manifestation due to excessive activation of tachykinergic neurons, and in particular excessive contraction of smooth muscle in any pathological condition in which release of tachykinins concurs in the genesis of the corresponding disorders.

In particular, the bronchospastic and inflammatory component of asthma, of
20 coughing, of pulmonary irritation, intestinal spasms, spasms of the biliary tract, and local spasms of the bladder and ureter in the course of cystitis and kidney infections and colics may be considered conditions in which the administration of the compounds of formula (I), as NK-2 antagonists, may prove effective.

The use as anxiolytic agents should also be considered for those compounds
25 that are provided with the appropriate chemico-physical characteristics for penetration into the CNS.

The compounds of formula (I) that are the subject of the present invention are suited for administration for therapeutic purposes to higher animals and man through the parenteral, oral, inhalational and sublingual routes, achieving
30 pharmacological effects according to the properties described above. For administration through parenteral (intravenous, intramuscular, and intradermal)

inhalational and sublingual instillation routes are concerned, aqueous solutions, aerosol preparations, powders or capsules are used according to the particular case.

The doses of active principle in the aforesaid compositions may range between
5 0.02 and 10 mg/kg of body weight.

The invention will now be illustrated in the examples that follow, which, however, have no limiting effect.

Example 1

Cyclo{-Suc-Trp-Phe-[(R)-NH-CH(CH₂C₆H₅)-CH₂-NH-]}

10 (compound of formula (I) where: X₁ = X₂ = X₃ = X₄ = -CO-NH-; R₁ = -CH₂-(indol-3-yl); R₂ = R₃ = -CH₂-C₆H₅; R₄ = R₅ = R₆ = R₇ = H; m = h = 0, f = g = 1; the carbon atoms C-R₁ and C-R₂ have an S configuration, whereas C-R₃ has an R configuration)

a) Synthesis of BOC-Trp-Phe-OH dipeptide

15 Di-*tert*-butyl carbonate (3.4 g) was added to a solution of H-Trp-Phe-OH (5 g) in dioxane (30 ml), H₂O (15 ml) and NaOH 1M (15.6 ml), cooled to 0-5°C under stirring. The reaction mixture was kept stirred for 2 hours, and then concentrated and extracted with pentane (2 x 20 ml). The aqueous phase was cooled with ice, with the addition of AcOEt (50 ml), KHSO₄ to obtain pH 2-3,
20 separated and extracted with AcOEt (2 x 50 ml). The re-united organic phases were washed with brine (50 ml), vacuum dried and evaporated at 30°C to obtain 6 g of the desired compound as a white semi-solid residue.

TLC: r.f. 0.55 (chloroform / cyclohexane / AcOH / H₂O = 45 / 45 / 5 / 5), 0.52 .
(CHCl₃ / MeOH = 9/1)

25 b) Synthesis of (R)-1-benzyl-2-benzyloxycarbonylamino-ethyl amine

The synthesis was carried out following the method described by P.G. Mattingly, Synthesis, 1990, 366, starting from BOC-D-phenylalaninol

c) Synthesis of BOC-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂-NH-Z] (5)

(R)-1-benzyl-2-benzyloxycarbonylamino ethyl amine (750 mg), PyBOP (1.37 g),
30 and DIEA (0.9 ml) were added to a solution of BOC-Trp-Phe-OH (1.19 g, 2.63 mmol.) in anhydrous DMG (10 ml) under nitrogen.

The reaction mixture was kept stirred overnight at room temperature, AcOEt (80 ml) was added, and the mixture was washed with HCl 1N (3 x 30 ml), Na₂CO₃ 5% (3 x 30 ml), and H₂O (30 ml). The organic phase was vacuum evaporated at 30°C to obtain 1.8 g of an ivory-coloured solid residue.

- 5 The crude compound was purified by washing in suspension with AcOEt under heat and with MeOH at room temperature to obtain 1.15 g of the desired product 5 as a white solid. MS(TS) : [MH⁺] = 718

d) Synthesis of H-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂-NH-Z] (6)

- TFA (6 ml) was added, under stirring and at room temperature, to a suspension
10 of the compound 5 (1.1 g) in CHCl₃ (30 ml), and a clear solution was seen to form immediately. The reaction mixture was kept stirred for 1.5 hours, and the disappearance of the precursor was monitored by means of HPLC analysis. After evaporation of the solvent, the residue was diluted with AcOEt (100 ml), washed with NaHCO₃ 5% (2 x 30 ml) and brine (30 ml).

- 15 The organic phase was dried with MgSO₄ and vacuum evaporated at 30°C.

The solid residue was purified by means of flash-chromatography (CHCl₃/MeOH = 95/5) to obtain 821 mg of the desired compound 6 as a white solid.

TLC: r.f. 0.50 (CHCl₃/MeOH = 9/1)

- 20 e) Synthesis of HO-Suc-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂-NH-Z] (compound 8 where: PG₂ = OH; PG₁ = Z)

NEt₃ (0.095 ml) and succinic anhydride (68 mg) were added to a solution of compound 6 (420 mg) in anhydrous DMF (10 ml) under stirring and at room temperature. The reaction mixture was kept stirred at room temperature for 4
25 hours.

After evaporation of the solvent, the residue was suspended in H₂O and kept stirred for 5 minutes. The solid was filtered and washed in suspension twice using MeOH to obtain 242 mg of the desired compound 8 as a white solid.

TLC: r.f. 0.50 (CHCl₃/MeOH = 8/2)

- 30 f) Synthesis of HO-Suc-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂-NH₂] (9)

The compound 8 (225 mg) was suspended in MeOH (10 ml) and hydrogenated in the presence of Pd/C 10% (50 mg) at atmospheric pressure and room

temperature. HPLC analysis after 4 hours showed that the precursor had disappeared completely.

The catalyst was filtered and washed with MeOH. After evaporation of the solvent, 158 mg of the desired compound 9 were obtained as a white solid.

5 m.p. = 142-4°C; TLC: r.f. 0.70 (*n*-butanol / AcOH / H₂O = 6 / 2 / 2)

g) Synthesis of cyclo{Suc-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂-NH-]}(10)

PyBOP (145 mg) and DIEA (0.09 ml) were added to a solution of compound 9 (148 mg) in anhydrous DMF (5 ml) stirred under nitrogen.

10 The reaction mixture was kept stirred for 5 hours and, after evaporation of the solvent, the residue was suspended in AcOEt, kept stirred for 10 minutes, and filtered, to obtain 100 mg of a solid product.

Part of the product (50 mg) was purified by HPLC to obtain 18 mg of the desired compound 10 as a white solid.

15 MS (TS) : [MH⁺] = 566; 1H-NMR (DMSO): d 2.15-2.35 (m, 2H), 2.55-2.85 (m, 8H), 2.96-3.04 (m, 2H), 3.90-4.02 (m, 1H), 4.03-4.15 (m, 1H), 4.25-4.42 (m, 1H), 6.71 (d, 1H), 6.90-7.42 (m, 16H), 8.09 (m, 1H), 8.50 (d, 1H), 10.82 (s, 1H).

Following the procedure described in Example 1, the compounds specified below were obtained:

Example 2

20 Cyclo{-Suc-Trp-Phe-[(S)-NH-CH(CH₂C₆H₅)-CH₂-NH-]}

(compound of formula I in which the substituents are defined as in Example 1, but all the C-R₁, C-R₂ and C-R₃ atoms have an S configuration) 1H-NMR (DMSO): d 1.95-2.32 (m, 2H), 2.34-2.90 (m, 6H), 2.92-3.18 (m, 2H), 3.60-3.82 (m, 1H), 4.00-4.40 (m, 4H), 6.90-7.36 (m, 14H), 7.39-7.54 (m, 2H), 7.64 (d, 1H),
25 7.88 (t, 1H), 8.27 (d, 1H), 10.78 (s, 1H).

Example 3

Cyclo{-Suc-Trp-Phe-[(R)-NH-CH(CH₂C₆H₁₁)-CH₂-NH-]}

(compound of formula I in which X₁ = X₂ = X₃ = X₄ = CO-NH-; R₁ = -CH₂-(indol-3-yl); R₂ = -CH₂-C₆H₅; R₃ = -CH₂-C₆H₁₁; R₄ = R₅ = R₇ = H; m = h = 0, f = g = 1;
30 the carbon atoms C-R₁ and C-R₂ have an S configuration, whereas C-R₃ has an R configuration) 1H-NMR (DMSO): d 0.65-0.95 (m 2H), 1.00-1.38 (m, 6H), 1.45-1.75 (m, 5H), 2.05-2.30 (m, 2H), 2.40-2.85 (m, 6H), 3.20-3.60 (m, 1H), 3.61-

3.78 (m, 1H), 3.80-3.96 (m, 1H), 3.98-4.10 (m, 1H), 4.38-4.55 (m, 1H), 8.47 (d, 1H), 6.90-7.45 (m, 11H), 8.02 (m, 1H), 8.47 (d, 1H), 10.78 (d, 1H).

Example 4

Cyclo{-Suc-Trp-Phe-[(R)-NH-CH(CH₂C₆H₄(4-OCH₃))-CH₂-NH-]}

- 5 (compound of formula I in which X₁ = X₂ = X₃ = X₄ = CO-NH-; R₁ = -CH₂-(indol-3-yl); R₂ = -CH₂-C₆H₅; R₃ = -CH₂-C₆H₄(4-OCH₃); R₄ = R₅ = R₆ = R₇ = H; m = h = 0, f = g = 1; the carbon atoms C-R₁ and C-R₂ have an S configuration, whereas C-R₃ has an R configuration) 1H-NMR (DMSO): d 2.13-2.37 (m, 2H), 2.50-2.85 (m, 8H), 3.25-3.50 (m, 1H), 3.58-3.80 (m, 4H), 3.85-4.00 (m, 1H), 4.02-4.18 (m, 10H), 4.28-4.45 (m, 1H), 6.65-7.47 (m, 16H), 8.02-8.16 (m, 1H), 8.48 (d, 1H), 10.80 (s, 1H).

Example 5

Cyclo{-Suc-Trp(5F)-Phe-[(R)-NH-CH(CH₂C₆H₅)-CH₂-NH-]}

- 15 (compound of formula I, in which R₁ = -CH₂-(5-fluoroindol-3-yl), and the other substituents are as defined in Example 1) MS(ES⁺): [MH⁺]=584

Example 6

Cyclo{-Suc-Trp(Me)-Phe-[(R)-NH-CH(CH₂C₆H₅)-CH₂-NH-]}

- (compound of formula I, in which R₁ = -CH₂-(N-methylindol-3-yl), and the other substituents are as defined in Example 1) MS(ES⁺):[MH⁺]= 580

20 Example 7

Cyclo{-Suc-Phe(3,4-Cl)-Phe-[(R)-NH-CH(CH₂C₆H₅)-CH₂-NH-]}

- (compound of formula I, in which R₁ = -(3,4-dichlorobenzyl), and the other substituents are as defined in Example 1) MS(ES⁺):[MH⁺]=595

Example 8

25 Cyclo{-Suc-Trp-Phe(3,4-Cl)-[(R)-NH-CH(CH₂C₆H₅)-CH₂-NH-]}

- (compound of formula I, in which R₂ = -(3,4-dichlorobenzyl), and the other substituents are as defined in Example 1) (MS(ES⁺):[MH⁺]= 634

Example 9

Cyclo{-Suc-Trp-Tyr-[(R)-NH-CH(CH₂C₆H₅)-CH₂-NH-]}

- 30 (compound of formula I, in which R₂ = -(4-hydroxybenzyl), and the other substituents are as defined in Example 1) (MS(ES⁺):[MH⁺]= 582

Cyclo{-Suc-Trp-Phe-[(R)-NH-CH(CH₂C₆H₃-3,4-diCl)-CH₂-NH-]}

(compound of formula I, in which R₃ = -(3,4-dichlorobenzyl), and the other substituents are as defined in Example 1) (MS(ES+):[MH+]= 634

Example 11

5 Cyclo{-Suc-Trp-Phe-[(R)-NH-CH(CH₂C₆H₄-4-OH)-CH₂-NH-]}

(compound of formula I, in which R₃ = -(4-hydroxybenzyl), and the other substituents are as defined in Example 1) (MS(ES+):[MH+]= 582

Example 12

Cyclo{-Suc-Trp-Phe-[(R)-NH-CH(CH₂-CH₂-C₆H₅)-CH₂-NH-]}

10 (compound of formula I, in which R₃ = -CH₂-CH₂-C₆H₅, and the other substituents are as defined in Example 1) (MS(ES+):[MH+]= 580

Example 13

Cyclo{-Suc-Trp-Phe-[(R)-NH-CH(CH₂-2-naphthyl)-CH₂-NH-]}

15 (compound of formula I, in which R₃ = -CH₂-(2-naphthyl), and the other substituents are as defined in Example 1) (MS(ES+):[MH+]= 616

Example 14

Cyclo{-Suc-Trp-Phe-[(R)-NH-CH(CH₂-indol-3-yl)-CH₂-NH-]}

(compound of formula I, in which R₃ = -CH₂-(indol-3-yl), and the other substituents are as defined in Example 1) (MS(ES+):[MH+]= 605

20 Example 15

Cyclo{-Suc-Trp-Phe-[(R)-NH-CH(CH₂-5-F-indol-3-yl)-CH₂-NH-]}

(compound of formula I, in which R₃ = -CH₂-(5-fluoroindol-3-yl), and the other substituents are as defined in Example 1) (MS(ES+):[MH+]= 623

Example 16

25 Cyclo{-Suc-Trp-Phe-[(R)-NH-CH(CH₂C₆H₄-3-F)-CH₂-NH-]}

(compound of formula I, in which R₃ = -CH₂C₆H₄-3-F, and the other substituents are as defined in Example 1) (MS(ES+):[MH+]= 584

Example 17

Cyclo{-Suc-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₃-3,4-diF)-CH₂-NH-]}

30 (compound of formula (I) wherein R₃ = -(3,4-difluorobenzyl) and the other substituents are as defined in Example 1 MS (ES+): [MH+]= 602

Example 18

Cyclo{-Suc-Trp-Phe-[(R)NH-CH(CH₂-C₆H₄-4-CF₃-CH₂-NH)-]}

(compound of formula (I) wherein R₃ = -(4-trifluoromethylbenzyl) and the other substituents are as defined in Example 1) MS (ES⁺): [MH⁺] = 634

Example 19

5 Cyclo{-Suc-Trp-Phe-[(R)-NH-CH₂-CH(CH₂C₆H₅)-NH-]}

(compound of formula (I) where: X₁ = X₂ = X₃ = X₄ = -CO-NH-; R₁ = -CH₂-(indol-3-yl); R₂ = R₃ = -CH₂-C₆H₅; R₄ = R₅ = R₆ = R₇ = H; f = h = 0; m = g = 1; the carbon atoms C-R₁ and C-R₂ have an S configuration, whereas C-R₃ has an R configuration)

10 a) Synthesis of (R)-2-*tert*-butoxycarbonylamino-3-phenyl-propylamine

The synthesis was performed according to the method described by P.G. Mattingly, Synthesis, 1990, 366, starting from BOC-D-phenylalaninol.

b) Synthesis of Z-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-NH-BOC] (5)

(R)-2-*tert*-butoxycarbonylamino-3-phenyl-propylamine (titre 65%, 1.1 g), PyBOP (1.45 g), and DIEA (0.98 ml) were added to a solution of Z-Trp-Phe-OH (1.4 g) 15 in anhydrous DMF (15 ml) under nitrogen. The reaction mixture was kept stirred overnight at room temperature, AcOEt (100 ml) was added, and the mixture was washed with HCl 1N (3 x 30 ml), Na₂CO₃ 5% (3 x 30 ml), and H₂O (30 ml). During the washings, the product partly precipitated, and was filtered and re- 20 united to the organic phase. After vacuum evaporation of the solvent, the residue (2.4 g) was washed in suspension with AcOEt and vacuum dried on P₂O₅, to obtain 1.79 g of the desired compound 5 as a white solid.

TLC: r.f. 0.86 (CHCl₃/MeOH = 95/5); r.f. 0.78 (AcOEt)

c) Synthesis of H-Trp-Phe-[(R)-NH-CH₂-CH(CH₂-C₆H₅)-NH-BOC] (6)

25 A suspension of the compound 5 (1.7 g) in MeOH (350 ml) was hydrogenated in the presence of Pd/C 10%, at atmospheric pressure and room temperature, until the precursor disappeared (HPLC analysis). After elimination of the catalyst by filtration and vacuum evaporation of the solvent, the residue was

d) Synthesis of HO-Suc-Trp-Phe-[(R)-NH-CH₂-CH(CH₂-C₆H₅)-NH-BOC]
(compound 8 where: PG₂ = OH; PG₁ = BOC)

Succinic anhydride (158 mg) and NEt₃ (0.21 ml) were added to a solution of compound 6 (840 mg) in anhydrous DMF (20 ml) under nitrogen. The reaction mixture was kept stirred at room temperature overnight. After vacuum evaporation of the solvent at a temperature of 30°C, the residue was treated with H₂O at 40-50°C, filtered, washed in suspension with MeOH (15 ml), and vacuum dried to obtain 600 mg of the desired compound 8 as a white solid.

TLC: 0.63 (CHCl₃/MeOH = 8/2)

e) Synthesis of HO-Suc-Trp-Phe-[(R)-NH-CH₂-CH(CH₂-C₆H₅)-NH₂]• TFA (9 TFA)

TFA (2ml) was added, under stirring, to a suspension of compound 8 (560 mg) in CH₂Cl₂ (15 ml), and a clear solution was obtained. After 2 hours at room temperature, the solvent was evaporated, and the residue diluted with ether, filtered and dried to obtain 500 mg of the desired compound 9 TFA as an ivory-coloured solid.

TLC: 0.58 (CHCl₃/MeOH = 8/2), 0.74 (*n*-butanol/AcOH/H₂O = 6/2/2)

f) Synthesis of cyclo{-Suc-Trp-Phe-[(R)-NH-CH₂-CH(CH₂-C₆H₅)-NH-]} (10)

PyBOP (447 mg), and DIEA (0.37 ml) were added, under nitrogen, to a solution of 9 TFA (500 mg) in anhydrous DMF (20 ml). The reaction mixture was kept stirred overnight at room temperature. After evaporation of the solvent, the residue was washed in suspension with citric acid 5% and H₂O. The product was dried on P₂O₅, washed in suspension using AcOEt and MeOH under heat, to obtain 110 mg of a solid. A portion was purified by HPLC to obtain 25 mg of the desired compound 10 as a white solid.

¹H-NMR (DMSO): δ 2.10-2.40 (m, 4H), 2.45-2.58 (m, 1H), 2.60-3.05 (m, 7H), 3.80-3.90 (m, 1H), 3.92-4.05 (m, 1H), 4.20-4.38 (m, 1H), 6.90-7.40 (m, 16H), 7.52-7.58 (m, 1H), 8.11 (d, 1H), 8.37 (d, 1H), 10.79 (s, 1H).

Example 20

Cyclo{-Suc-Trp-Phe-[(S)-NH-CH₂-CH(CH₂C₆H₅)-NH-]}

(compound of formula I in which the substituents are defined as in Example 19, except for the fact that C-R₃ has an S configuration).

The compound was obtained following a procedure similar to that described for Example 19.

¹H-NMR (DMSO): d 1.98-2.26 (m, 2H), 2.40-2.88 (m, 8H), 2.98-3.11 (m, 1H), 3.66-3.84 (m, 1H), 3.98-4.23 (m, 2H), 4.40-4.58 (m, 1H), 6.89-7.48 (m, 17H),
5 8.10 (d, 1H), 8.44 (d, 1H), 10.83 (s, 1H).

Proceeding in a similar way as that described in Example 1 above, the following compounds were obtained:

Example 21

Cyclo{-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂-NH-]-(CH₂)₃CO-}
10 (compound of formula I, in which R₃ = -CH₂-C₆H₅ and X₃ = -CH₂-NH-, and the other substituents are as defined in Example 1. MS (ES⁺): [MH⁺]=552.

Example 22

Cyclo{-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂-N(CH₃)]-(CH₂)₃CO-}
(compound of formula I, wherein R₃= -CH₂-C₆H₅ and X₃=-CH₂N(CH₃)- and the
15 other substituents are as defined in Example 1. MS(ES⁺):[MH⁺]=566.

EXAMPLE 23

Cyclo{-Suc[1(S)-NH₂]-Trp-Phe-[(R)NH-CH(CH₂-C₆H₅)-CH₂NH]-}
(compound of formula I wherein h = 1, g = 0, R₄ = -NH₂ and the other
substituents are as defined in Example 1 while the carbon atom C-R₄ has
20 configuration S).

a) Synthesis of Boc-Asp[Trp-Phe-[(R)NH-CH(CH₂-C₆H₅)-CH₂-NH-Z]-OBz]

(compound 8 wherein: PG₂ = OBzl, PG₁ = Z

To a solution of compound 6 (see Example 1d) (650 mg) in anhydrous DMF (30 ml) Boc-Asp-OBzl (340 mg), PyBOP (656 mg) and ET₃N (0.4 ml) are added
25 under stirring at room temperature. The mixture is stirred for 2 h at room temperature. After evaporation of the solvent under vacuum the residue was treated with H₂O giving a solid residue which is filtered, washed with water and

having a linear gradient of acetonitrile containing 0.1%(v/v) TFA (phase B) against aqueous TFA 0.1% (v/v) (phase A), from 20% to 80% in B in 20 minutes at a flux of 3 ml/min; determination by UV at 220 nm. Retention time: $R_t = 21.1$ min.

5 b) Synthesis of Boc-Asp[Trp-Phe-[(R)NH-CH(CH₂-C₆H₅)-CH₂-NH₂]]-OH (9)

The compound 8 (of Example 23a) (600 mg) was solved in DMF (2 ml) and diluted with MeOH (30 ml), hydrogenated in the presence of Pd/C 10% (100 mg) at room pressure and temperature for 5 h. The catalyser was filtered and washed with MeOH. After evaporation of the solvent 420 mg of the desired product 9 were obtained in the form of a white solid.

MS(ES⁺):[MH⁺]= 663; HPLC (same conditions as above): $R_t = 11.07$.

c) Synthesis of cyclo{-Suc[1(S)NH-BOC]-Trp-Phe[(R)NH-CH(CH₂-C₆H₅)-CH₂-NH]-} (10)

To a solution of compound 9 (see example 23b) (7.2 g) in anhydrous DMF (900 ml) 4 g of HBT and 2 g of EDC were added under stirring and nitrogen atmosphere. The mixture was stirred for 5 h and, after evaporation of the solvent, the residue was treated with an aqueous solution of KHSO₄ 5% and extracted in ethylacetate.

The organic phase was washed with brine, NaHCO₃ 5% and again with brine, dried and evaporated the yellow solid obtained (5.2 g) was crystallized from isopropanol/water: 1/1 giving 3.2 g of a white solid. MS(ES⁺):[MH⁺]=681; HPLC (same conditions as above): $R_t = 14.8$.

d) Synthesis of cyclo{-Suc[(1(S)NH₂)-Trp-Phe-[(R)NH-CH(CH₂-C₆H₅)-CH₂-NH]]} (10)

To a suspension of compound 10 (see example 23d) (1g) in CH₂Cl₂ (20 ml) TFA (7 ml) was added under stirring at 0°C giving a clear solution; thereafter the temperature is raised up to room temperature. The mixture was left at room temperature for 90 minutes and then the solvent was evaporated and the residue was treated with NaHCO₃ and water and extracted in ethylacetate. The organic phase was washed with brine, dried and evaporated giving a solid (800 mg).

MS(ES⁺):[MH⁺]= 581; HPLC (same conditions as above said): $R_t = 9.4$.

A sample of 20 mg is purified by HPLC giving 15 mg of trifluoroacetate: cyclo{-Suc[1(S)NH₂]-Trp-Phe-[(R)NH-CH(CH₂-C₆H₅)-CH₂-NH]-}.TFA (10 TFA)

MS(ES⁺):[MH⁺]= 581; HPLC: Rt= 9.4 (same conditions as above); ¹H-NMR (DMSO): δ 2.60-2.90 (m, 8H), 3.05-3.11 (m, 1H), 3.63-3.71 (m, 1H), 4.07-4.13 (m, 3H), 4.32-4.38 (m, 1H), 6.90-7.45 (m, 17H), 8.07 (bs, NH₃⁺), 8.22-8.28 (m, 1H), 8.57 (d, 1H), 10.82 (s, 1H).

Following the procedure described in Example 23 the following compounds were obtained:

Example 24

10 Cyclo{-Suc[1(R)-NH₂]-Trp-Phe-[(R)NH-CH(CH₂-C₆H₅)-CH₂NH]-}

(compound of formula I wherein h = 1, g = 0, R₄ = -NH₂ and the other substituents are as defined in Example 1 while the carbon atom C-R₄ has configuration R) MS (ES⁺): [MH⁺] = 581

Example 25

15 Cyclo{-Suc[2(S)-NH₂]-Trp-Phe-[(R)NH-CH(CH₂-C₆H₅)-CH₂NH]-}

(compound of formula I wherein h = 1, g = 0, R₄ = -NH₂ and the other substituents are as defined in Example 1 while the carbon atom C-R₄ has configuration S) MS (ES⁺): [MH⁺] = 581

Example 26

20 Cyclo{-Suc[2(R)-NH₂]-Trp-Phe-[(R)NH-CH(CH₂-C₆H₅)-CH₂NH]-}

(compound of formula I wherein h = 1, g = 0, R₄ = -NH₂ and the other substituents are as defined in Example 1 while the carbon atom C-R₄ has configuration R) MS (ES⁺): [MH⁺] = 581

Example 27

25 Cyclo{-Suc[1(S)-NH(CH₃)]-Trp-Phe-[(R)NH-CH(CH₂-C₆H₅)-CH₂NH]-}

(compound of formula I wherein h = 1, g = 0, R₄ = -NH(CH₃) and the other substituents are as defined in Example 1 while the carbon atom C-R₄ has configuration S) MS (ES⁺): [MH⁺] = 595

(compound of formula I wherein $h = 1$, $g = 0$, $R_4 = -\text{COO}(\text{CH}_2-\text{C}_6\text{H}_4-4-\text{NO}_2)$ and the other substituents are as defined in example 23) (diastereoisomeric mixture in respect of C- R_4 and separation of the two epimers).

a) Synthesis of Boc-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂-NH₂]

- 5 The Boc-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂-NH-Z] (5 see example 1c) (1.2 g) was dissolved in a mixture of DMF (30 ml) and MeOH (200 ml) and hydrogenized in the presence of Pd/C 10% (200 mg) at room pressure and temperature, for 4 h. The catalyser was filtered and washed with MeOH, the solvent evaporated giving 700 mg of solid residue.

- 10 MS(ES⁺): [MH⁺] 584; HPLC (conditions of example 23): Rt = 11.1

b) Boc-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂-NH]-COCH[COO(CH₂-C₆H₄-4-NO₂)]CH₂COO-tBu

- 424 mg of 2-(4-nitro-benzyloxycarbonyl)-succinic acid 4-tert-butyl ester were dissolved in DMF (20 ml). To the mixture HOBt (490 mg), EDC and Boc-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂-NH₂] were added at 0°C under stirring; the temperature was raised to room temperature while stirring for 2 h. The solvent was evaporated and the residue treated with KHSO₄ 5% giving a yellow solid which was filtered, washed with NaHCO₃ 5%, water and dried.

- 1.05 g of compound were obtained, MS(ES⁺):[MH⁺] = 919; HPLC (conditions of Example 23): Rt = 20.36

c) H-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂-NH]-COCH[COO(CH₂-C₆H₄-4-NO₂)]CH₂COOH

- Boc-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂-NH]-COCH[COO(CH₂-C₆H₄-4-NO₂)]CH₂COO-tBu (1.05 g) was added in small portions in anhydrous trifluoroacetic acid (20 ml) at 0°C and the mixture was kept under stirring for 30 minutes, dried and the residue treated with ethyleter; the formed solid was filtered, washed with ethyleter and dried, 850 mg of product were obtained.

MS(ES⁺):[MH⁺] = 763; HPLC (conditions of example 23): Rt = 10.6

- d) Synthesis of cyclo{-Suc[1-COO(CH₂C₆H₄-4-NO₂)]-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂-NH-]}

The H-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂-NH]-COCH[COO(CH₂C₆H₄-4-NO₂)]CH₂COOH (100 mg) was dissolved in DMF (5 ml) and to the mixture PyBOP (80 mg) and Et₃N (54 μ l) were added stirring for 3 h.

The reaction mixture was dried and the residue dissolved in ethylacetate, the organic phase was washed with KHSO₄ 5%, brine, NaHCO₃ 5% and brine, dried and concentrated. 90 mg of epimeric mixture was obtained, the epimers were separated by HPLC giving:

30 mg of liophylized solid which in HPLC (conditions of example 23) shows an Rt = 15.2. MS(ES+):[MH+]= 745.

1H-NMR (DMSO): δ 2.54-2.81 (m, 7H), 3.08-3.17 (m, 1H), 3.34-3.39 (m, 1H), 3.77-3.84 (m, 1H), 4.00-4.10 (m, 3H), 4.37-4.43 (m, 1H), 5.31 (s, 2H), 6.60 (d, 1H), 6.93-7.42 (m, 16H), 7.62 (d, 2H), 8.24 (d, 2H), 8.60 (d, 1H), 8.66-8.72 (m, 1H), 10.81 (s, 1H) and

7 mg of liophylized solid which in HPLC (conditions of example 23) shows an Rt = 15.7. MS(ES+): [MH+]= 745.

Example 29

Cyclo{-Suc(1-COOH)-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂-NH]} (compound of formula I, wherein h = 1, g = 0, R₄ = -COOH and the other substituents are as defined in Example 1) [epimer which in HPLC (conditions of Example 23) shows an Rt = 10.7]

The cyclo{-Suc(1-COO(CH₂-C₆H₄-4-NO₂))-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂-NH]} which in HPLC (same conditions as in example 23) shows an Rt = 15.2 (50 mg) was suspended in a mixture water/isopropanol:1/1 (6 ml) containing K₂CO₃ (19 mg) and was kept under stirring for 24 h. The solvent was evaporated and the residue was diluted with water and the solution washed with ethylacetate, by adding HCl 1N separated a solid which was extracted with ethylacetate; the organic phase was washed with brine and dried. By evaporating the solvent mg 35 of a solid residue were obtained.

MS(ES+):[MH+]=610. HPLC (conditions of Example 23): Rt = 10.7

Example 30

Cyclo{-Suc(1-COOH)-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂-NH]} (compound of formula I, wherein h = 1, g = 0, R₄ = -COOH and the other substituents are as

defined in Example 1) [epimer which in HPLC (conditions as in Example 23) shows an $R_t = 11.1$]

The cyclo{-Suc(1-COO(CH₂-C₆H₄-4-NO₂))-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂-NH-]} having $R_t = 15.7$ was hydrolyzed as described in Example 29.

- 5 MS(ES⁺):[MH⁺] = 610; HPLC (same conditions of Example 23): $R_t = 11.1$
As described in Example 28 the following compounds were obtained :

Example 31

Cyclo{-Suc(1-OH)-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂-NH-]}

- (compound of formula I, in which $h = 1$; $g = 0$; $R_4 = -OH$, and the other
10 substituents are as defined in Example 1), MS(ES⁺):[MH⁺] = 582.

Example 32

Cyclo{-Suc(2-COOH)-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂-NH-]}

(compound of formula I, in which $R_4 = -COOH$, and the other substituents are
as defined in Example 1) MS(ES⁺):[MH⁺] = 610.

- 15 Example 33

Cyclo{-Suc(2-OH)-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂-NH-]} (compound of
formula I wherein: $h = 0$, $g = 1$, $R_4 = OH$ and the other substituents are as
defined in example 1) MS(ES⁺): [MH⁺] = 582.

- The compounds of Examples 23, 24, 25, 26, 27, 29, 30 and 32 can be
20 derivatized as described hereinafter.

Example 34

Cyclo{-Suc[1(S)-(2H-tetrazolyl-5-ylmethyl)amino]-Trp-Phe-[(R)-NH-CH(CH₂-
C₆H₅)-CH₂-NH]-}.TFA (compound of formula I wherein $h = 1$, $g = 0$, $R_4 = -(2H$ -
tetrazolyl-5-ylmethyl)amino and the other substituents are as defined in
25 example 1 while the carbon atom C- R_4 has configuration S)

a) Synthesis of 5-iodomethyl-1-trityl-1H-tetrazole

- To a suspension of 5-chloromethyl-1H-tetrazole (6.0 g) in chloroform (100 ml)
trityl-chloride (14.2 g) was added at 0°C under nitrogen, and the mixture was
stirred up to total solubilization, thereafter a solution of Et₃N (7.0 ml) in
30 chloroform (50 ml) was added at 5°C and the temperature was left raising up to
room temperature, the mixture was kept resting for 24 h.

The mixture was treated with ethylacetate (200 ml) and left resting for 6h, the separated solid was filtered away and to the solution acetone (70 ml) was added, the precipitated solid was collected by filtration and dried giving 9.5 g of 5-chloromethyl-1-trityl-1H-tetrazole which was solubilized in acetone (200 ml) at 60°C. Sodium iodide (5.6 g) was added to the solution which was refluxed for 6 h, by cooling precipitated a compound which was filtered, washed with water and dried giving 5.2 g of a white solid.

TLC: R.f. 0.55 (AcOEt/Cyclohexane : 1/3)

b) Synthesis of cyclo{-Suc[1(S)-(2-trityl-tetrazolyl-5-ylmethyl)amino]-Trp-Phe-[(R)NH-CH(CH₂-C₆H₅)-CH₂NH]-}

To 205 mg of cyclo{-Suc[1(S)-NH₂]-Trp-Phe-[(R)NH-CH(CH₂-C₆H₅)-CH₂NH]-} (compound of example 23) in anhydrous DMF (5 ml) were added, under stirring 5-iodomethyl-1-trityl-1H-tetrazole (147 mg) and thereafter DIEA (0.06 ml) keeping the temperature at 0°C for 4 h and at room temperature for 3 h. The mixture was treated with water and extracted with ethylacetate, the organic phase was washed with brine and dried. By evaporating the solvent a solid was obtained which was purified by column-chromatography eluting with AcOEt/MeOH = 95/5. 210 mg of product were obtained. MS(ES⁺):[MH⁺]=905; HPLC (conditions of example 23): Rt=15.4.

c) synthesis of cyclo{-Suc[1(S)-(2-HI-tetrazolyl-5-ylmethyl)amino]-Trp-Phe-[(R)NH-CH(CH₂-C₆H₅)-CH₂NH]-}.TFA

To a solution of cyclo{-Suc[1(S)-(2-trityl-tetrazolyl-5-ylmethyl)amino]Trp-Phe-[(R)NH-CH(CH₂-C₆H₅)-CH₂NH]-} (90 mg) in anhydrous DMF (5 ml) a solution of HCl 4M in dioxane (0.6 ml) was added at 0°-5°C, the temperature was brought to room temperature and the mixture was left resting up to end of the reaction (14 h at room temperature and 56 h at 5°C) checking the reaction by HPLC

¹H-NMR (DMSO): δ 2.62-2.92 (m, 8H), 3.16-3.23 (m, 1H), 3.68-3.74 (m, 1H), 4.00-4.14 (m, 3H), 4.25-4.75 (m, 3H), 6.88-7.42 (m, 17H), 8.30-8.37 (m, 1H), 8.54 (d, 1H), 10.82 (s, 1H).

Example 35

- 5 Cyclo{-Suc[1(S)-(morpholin-4-yl)]-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂-NH]-}.TFA (compound of formula I wherein h = 1, g = 0, R₄ = -(morpholin-4-yl) and the other substituents are as defined in example 1 while the carbon atom C-R₄ has configuration S)

To a solution of 2.2'-oxydiacetaldehyde (1mmole), excess, in methanole (20 ml)
10 58 mg of cyclo{-Suc[1(S)-NH₂]-Trp-Phe-[(R)NH-CH(CH₂-C₆H₅)-CH₂NH]-} (compound of example 23), 0.2 ml of acetic acid and 12 mg of NaCNBH₃ were added. After 2 h the mixture was diluted with water (10 ml), treated with HCl 1N up to pH 3 and the methanole was evaporated; the solution was treated with NaHCO₃ 5% and the formed solid was extracted with ethylacetate. The organic
15 phase, after washing with brine and anhydrification, was evaporated giving 58 mg of a solid which was purified by HPLC giving 10 mg of liophylized solid trifluoroacetate.

MS(ES⁺):[MH⁺]=651; TLC: R.f.0.20 (CHCl₃/MeOH:9/1)

¹H-NMR (DMSO): δ 2.62-3.00 (m, 8H), 3.27-3.87 (m, 10H), 4.07-4.15 (m, 3H),
20 4.32-4.38 (m, 1H), 6.62 (d, 1H), 6.94-7.41 (m, 16H), 8.49 -8.64(m, 2H), 10.84 (s, 1H).

Via a similar reductive amination reaction, as described in example 35, the following compounds were obtained:

Example 36

- 25 Cyclo{-Suc[1(S)-N(CH₃)₂]-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂NH]-}.TFA (compound of formula I wherein h = 1, g = 0, R₄ = -N(CH₃)₂ and the other substituents are as defined in example 1 while the carbon atom C-R₄ has configuration S)

The synthesis was performed starting from the compound of example 23 using
30 paraformaldehyde. MS(ES⁺):[MH⁺]=609.

Example 37

Cyclo{-Suc[1(S)-(piperidin-4-yl)]-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂NH]-}.TFA (compound of formula I wherein h = 1, g = 0, R₄ = -(piperidin-4-yl) and the other substituents are as defined in example 1 while the carbon atom C-R₄ has configuration S)

- 5 The synthesis was performed starting from the compound of example 23 using glutaraldehyde. MS(ES+):[MH+]=649.

Example 38

- Cyclo{-Suc[1(S)-(N(CH₂CH₂OH)₂)]-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂NH]-}.TFA (compound of formula I wherein h = 1, g = 0, R₄ = -N(CH₂CH₂OH)₂ and
10 the other substituents are as defined in example 1 while the carbon atom C-R₄ has configuration S)

The synthesis was performed starting from the compound of example 23 using glycolaldehyde. MS(ES+):[MH+]=669.

Example 39

- 15 Cyclo{-Suc[1(S)-(NHCH₂CH(OH)CH₂OH)]-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂-NH]-}.TFA (compound of formula I wherein h = 1, g = 0, R₄ = -NHCH₂CH(OH)CH₂OH and the other substituents are as defined in example 1 while the carbon atom C-R₄ has configuration S)

- The synthesis was performed starting from the compound of example 24 using
20 D-glyceraldehyde. MS(ES+):[MH+]=655.

Example 40

- Cyclo{-Suc[1(S)-(3-carboxypropanoyl)amino]-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂-NH]-}. (compound of formula I wherein h = 1, g = 0, R₄ = -(3-carboxypropanoyl)amino and the other substituents are as defined in example
25 1 while the carbon atom C-R₄ has configuration S)

- To a solution of the compound of Example 23 (100 mg) in anhydrous DMF (2 ml) succinic anhydride (30 mg) and dimethylamino-pyridine (10 mg) were added and the solution was stirred for 16 h; the solvent was evaporated giving a solid which was solubilized in ethylacetate, washed with citric acid 10%, brine and
30 dried. By evaporating the solvent a solid compound was collected (90 mg), which purified by HPLC gave 60 mg of a lipophilized solid.

MS(ES+):[MH+]=681; HPLC (conditions as in example 23): Rt=10.8

¹H-NMR (DMSO): δ 2.35-2.82 (m, 12H), 3.25-3.28 (m, 1H), 3.66-3.73 (m, 1H), 3.98-4.12 (m, 2H), 4.33-4.38 (m, 1H), 4.67-4.73 (m, 1H), 6.80 (d, 1H), 6.96-7.39 (m, 16H), 8.16-8.23 (m, 2H), 8.51 (d, 1H), 10.89 (s, 1H).

Example 41

5 Cyclo{-Suc[1(S)-[3-N'-(β -D-glucopiranos-1-yl)-carboxamidopropanoyl]amino]-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂-NH]-} (compound of formula I wherein h = 1, g = 0, R₄ = -[3-N'-(β -D-glucopiranos-1-yl)carboxyamidopropanoyl]amine] and the other substituents are as defined in example 1 while the carbon atom C-R₄ has configuration S)

10 The compound of example 40 (90 mg) was dissolved in anhydrous DMF (10 ml) under stirring and in nitrogen atmosphere, to the mixture 55 mg HBT, 25 mg EDC and 24 mg β -D-glucopiranosylamine were added.

The mixture was left stirring overnight and after evaporation of the solvent the resulting oil was treated with citric acid 10% giving a solid which was filtered,
15 washed with water and dried. The 80 mg obtained were purified by HPLC giving 40 mg of a lyophilized solid.

MS(ES⁺):[MH⁺]= 842; HPLC (conditions of example 23): Rt= 8.2

¹H-NMR (DMSO): δ 2.31-2.81 (m, 12H), 3.00-3.10 (m, 2H), 3.13-3.65 (m, 5H), 3.66-3.75 (m, 1H), 3.97-4.12 (m, 2H), 4.29-4.36 (m, 1H), 4.65-4.75 (m, 2H), 6.78
20 (d, 1H), 6.95-7.40 (m, 16H), 8.19-8.27 (m, 2H), 8.35 (d, 1H), 8.51 (d, 1H), 10.82 (s, 1H).

Example 42

Cyclo{-Suc[1(S)-[(carboxymethyl)amino]-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂-NH]-} TFA (compound of formula I wherein h = 1, g = 0, R₄ = -
25 (carboxymethyl)amino and the other substituents are as defined in example 1 while the carbon atom C-R₄ has configuration S)

a) Synthesis of cyclo{-Suc[1(S)-[(ter-butoxycarbonylmethyl)-amino]-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂-NH]-}

To a solution of the compound of example 23 (130 mg) in anhydrous DMF (3
30 ml) DIEA (0.04 ml) and ter-butyle (0.04 ml) bromoacetate were added, the solution was stirred for 2 h and thereafter the mixture was poured in KHSO₄

5%. The formed solid was filtered, washed with NaHCO_3 , water and dried. 100 mg of product were obtained.

HPLC (conditions of Example 23): $R_t = 11.3$

b) Synthesis of cyclo{-Suc[1(S)-[(carboxymethyl)amino]-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂-NH]-} .TFA

The above collected solid (90 mg) was suspended in CH_2Cl_2 (5 ml) and TFA (5 ml) was added under stirring at 0°C, the mixture was stirred for 1 h at room temperature. The solution was concentrated and the obtained residue was purified by HPLC giving 40 mg of a liophylized solid.

MS(ES+):MH+=639; HPLC (conditions of Example 23): $R_t=9.4$.

Example 43

Cyclo{-Suc[1(S)-[N'-(β-D-glucopiranos-1-yl)-carboxyamidomethyl]amino]-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂-NH]-} TFA (compound of formula I wherein h = 1, g = 0, R₄ = -[N'-(β-D-glucopiranos-1-yl)carboxyamidomethyl]amine] and the other substituents are as defined in example 1 while the carbon atom C-R₄ has configuration S)

The product was obtained starting from the product of Example 42 and β-D-glucopiranosylamine according to the procedure of Example 41.

MS(ES+):[MH+]=800; HPLC (conditions of example 23): $R_t= 7.6$

Example 44

cyclo{-Suc[1(S)-(chiny)amine]-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂-NH]-} (compound of formula I wherein h = 1, g = 0, R₄ = -(chiny)amine and the other substituents are as defined in example 1. while the carbon atom C-R₄ has configuration S)

Chinic acid (50 mg) was solubilized in anhydrous DMF (10 ml) under stirring and nitrogen atmosphere, HBT (220 mg), EDC (100 mg) and the compound obtained in Example 24 (150 mg) were added. The mixture was left under stirring overnight, thereafter the solvent was evaporated and the residue treated with an aqueous solution of KHSO_4 5% and extracted with ethylacetate.

The organic phase was washed with brine, NaHCO_3 5% and again brine, dried and evaporated; the obtained solid (122 mg) was purified on flash

chromatography (SiO₂) eluting with chloroform/methanol:8/2; 80 mg of the desired compound were obtained.

MS(ES⁺):[MH⁺]=755; HPLC (conditions as in example 23) Rt=10.05

Example 45

- 5 Cyclo{-Suc[1(S)-(4-aminobutanoyl)amino]-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂-NH]-} TFA (compound of formula I wherein h = 1, g = 0, R₄ = -(4-aminobutanoyl)amino and the other substituents are as defined in example 1 while the carbon atom C-R₄ has configuration S)

The product was obtained starting from the product of Example 23 and 4-BOC-aminobutirryc acid according to the procedure of Example 44 followed by
10 elimination of the protecting group BOC.

MS(ES⁺):[MH⁺]=666.

Example 46

- 15 Cyclo{-Suc[1(S)-[(1,4')bipiperidin-1-yl]acetamido]-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂-NH]-} TFA (compound of formula I wherein h = 1, g = 0, R₄ = -[(1,4')bipiperidin-1-yl]acetamido and the other substituents are as defined in example 1 while the carbon atom C-R₄ has configuration S)

The product was obtained starting from the product of Example 23 and [(1,4')bipiperidin-1-yl]acetic acid according to the procedure of Example 44.

20 MS(ES⁺):[MH⁺]=789.

Example 47

- 25 Cyclo{-Suc[1-N-(β-D-glucopiranos-1-yl)-carboxyamido]-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂-NH]-} (compound of formula I wherein h = 1, g = 0, R₄ = N-(β-D-glucopiranos-1-yl)carboxyamide and the other substituents are as defined in example 1 while the carbon atom C-R₄ has configuration S)

The product was obtained starting from the product of Example 29 and β-D-glucopiranosylamine according to the procedure of Example 44.

MS(ES⁺):[MH⁺]= 771.

Example 48

- 30 Cyclo{-Suc[1(S)-[N'-(2-N-acetyl-β-D-glucopiranos-1-yl)-carboxyamido]-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂-NH]-} (compound of formula I wherein h = 1, g = 0, R₄ = -N'-(2-N-acetyl-β-D-glucopiranos-1-yl)carboxyamide and the other

substituents are as defined in example 1 while the carbon atom C-R₄ has configuration S)

The product was obtained starting from the acid of Example 29 and 2-N-acetyl- β -D-glucopiranosylamine according to the procedure of Example 44.

5 MS(ES⁺):[MH⁺]=812.

Biological activity

The compounds described in the present invention act as antagonists of the NK-2 receptor of tachykinins. The biological activity was evaluated in two *in-vitro* functional tests, using rabbit pulmonary artery (RPA) and hamster trachea
10 (HT), according to the methods described by C.A. Maggi *et al.*, Br. J. Pharmacol., 1990, 100, 588 and P. D'Orléans-Juste *et al.*, Eur. J. Pharmacol., 1986, 125, 37. The activity of the compounds as human NK-2 receptor antagonists was assessed in a binding test using membranes of Chinese hamster ovary (CHO) cells, transfected with the NK-2 receptor of human ileum
15 and the radioligand [¹²⁵I]NKA (Amersham, specific activity 2000 Ci/mmol) at a concentration of 100 pM in competition studies. The substances under examination were tested in a concentration range of from 0.01 nM to 10 mM. At the end of incubation (30 minutes at 20°C), the samples were filtered on Whatman GF/B filters and employing the Brandel automatic filtration system.
20 Radio-activity was determined by means of a gamma counter (Cobra, Canberra Packard).

The data gathered from the functional studies were expressed as pA₂ (O. Arunlakshana and H.O. Schild, Br. J. Pharmacol. Chemother., 1959, 14, 48), and those of the binding studies as pKi (-log Ki calculated using the LIGAND
25 programme: P.J. Munson *et al.*, Anal. Biochem., 1980, 107, 220).

The compounds of the invention proved active in the tests referred to above, with pA₂ values of between 5 and 9, the more powerful compounds revealing a higher affinity for the human receptor, with pKi of between 8 and 10.

List of abbreviations used

30 For the nomenclature and abbreviations of amino acids, reference is made to the

Nomenclature (Eur. J. Biochem., 1984, 138, 9); the amino acids are understood in the S configuration, if not otherwise specified.

The other abbreviations used are the following:


BOC = *tert*-butoxycarbonyl; Z = benzyloxycarbonyl; -Suc- = succinyl; Bzl =
5 benzyl; PyBOP = (benzotriazol-1-yloxy)*tris*(pyrrolidine) phosphonium
hexafluorophosphate, DIEA = N,N-diisopropylethylamine; NEt₃ = triethylamine;
DMF = N,N-dimethylformamide; NKA = neurochinine A; TFA = trifluoro-acetic
acid; HBT = 1-hydroxybenzotriazole; EDC = N-(3-dimethylaminopropyl)-N'-
ethylcarbodiimide hydrochloride.

10 The numeration of the substituents on the succinic-group is as follows:

-Suc(1-NH₂)- = -CO-CH(NH₂)-CH₂-CO-

-Suc(2-NH₂)- = -CO-CH₂-CH(NH₂)-CO-


Scheme



where PG, PG1 and PG2 are protecting groups commonly used in the synthesis of peptides

CLAIMS

1. Monocyclic compounds having the general formula (I):



in which:

X_1, X_2, X_3, X_4 , which may be the same or different from one another, represent a group chosen from among $-CONR-$, $-NRCO-$, $-OCO-$, $-COO-$, $-CH_2NR-$, $-NR-CH_2-$, CH_2-CH_2 , where R is H or a C_{1-3} alkyl or benzyl;

f, g, h, m, which may be the same or different from one another, represent a number chosen from among 0, 1 or 2;

R_1 and R_2 , which may be the same or different from one another, represent a $-(CH_2)_r-Ar$ group, where $r = 0, 1, 2$ and where Ar is an aromatic group chosen from among: benzene, naphthalene, thiophene, benzothiophene, pyridine, quinoline, indole, furan, benzofuran, thiazole, benzothiazole, imidazole, and benzo-imidazole, the said Ar group being possibly substituted with a maximum of 2 residues chosen from among C_{1-3} alkyl or halo-alkyl, C_{1-3} alkoxy, C_{2-4} amino-alkoxy, halogen, OH, NH_2 , $NR_{13}R_{14}$ where R_{13} and R_{14} , which may be the same or different from one another, represent hydrogen or C_{1-3} alkyl;

R_3 represents a group chosen from among:

- hydrogen

- linear or branched alkyl having the formula C_nH_{2n+1} , with $n = 1-5$, cyclo-alkyl or alkylcyclo-alkyl groups having the formula C_nH_{2n-1} with $n = 5-9$

- $(CH_2)_r-Ar_1$, where $r = 0, 1, 2$ and where Ar_1 is an aromatic group chosen from among: benzene, naphthalene, thiophene, benzothiophene, pyridine, quinoline, indole, furan, benzofuran, thiazole, benzothiazole, imidazole, and benzo-imidazole, the said Ar_1 group being possibly substituted with a maximum of 2 residues chosen from among C_{1-3} alkyl or halo-alkyl, C_{1-3} alkoxy or amino-alkoxy, halogen, OH, NH_2 , $NR_{13}R_{14}$, where R_{13} and R_{14} , which may be the same or different from one another, represent hydrogen or C_{1-3} alkyl;

32 R_4 represents a group chosen from among:

33 - hydrogen or C_{1-6} alkyl

34 - L-Q, where L is a chemical bond or a linear or branched C_{1-6} alkyl residue and
35 Q is a group chosen from among:

36 i) H, OH, OR_9 , NH_2 , NR_9R_{10} , guanidine, sulphate, phosphonate, phosphate,
37 where R_9 and R_{10} , which may be the same or different from one another,
38 represent a hydrogen, C_{1-3} alkyl group, C_{1-3} hydroxyalkyl, C_{1-3} dihydroxyalkyl, C_{1-3}
39 alkyl- $CONHR_{12}$, C_{1-3} alkyltetrazole, C_{1-3} alkyl-COOH or wherein R_9R_{10} joined
40 together form with the N-atom a saturated 4-6 membered heterocycle possibly
41 containing a further heteroatom chosen in the group consisting of N, O, S and
42 wherein R_{12} is a mono-, di-, tri-glycosidic group possibly protected with one or
43 more C_{1-3} -acyl groups or substituted with amino-groups or C_{1-3} acylamino-
44 groups;

45 ii) COOH, tetrazole, SO_2NH_2 , $SO_2NHCOOR_8$, $CONHR_8$, $NHCOR_8$, where R_8
46 represents a linear or cyclic C_{1-6} alkyl chain containing one or more polar groups
47 chosen from among the group: OH, NH_2 , $NR_{15}R_{16}$, COOH, $CONHR_{12}$, PO_3H ,
48 SO_3H , OR_{11} and where R_{15} and R_{16} , which may be the same or different from
49 one another, represent a hydrogen or C_{1-3} alkyl group, and where R_{11} is a C_{1-3}
50 alkyl or C_{2-4} amino-alkyl chain, R_{12} is a mono-, di-, tri-glycosidic group possibly
51 protected with one or more C_{1-3} acyl groups or substituted with amino-groups or
52 C_{1-3} acylamino-groups or $R_{15}R_{16}$ joined together form with the N-atom a
53 saturated 4-6 membered heterocycle possibly substituted with C_{1-3} alkyl-groups
54 or with saturated 4-6 membered heterocycle-groups containing at least an N-
55 atom;

56 iii) $COOR_{17}$, $CONHR_{12}$, OR_{12} where R_{12} is a mono-, di- or tri-glycoside group
57 possibly protected with one or more C_{1-3} acyl groups or substituted with amine
58 or C_{1-3} acylamine groups and R_{17} is a group R_{12} as above defined or a group
59 C_{1-3} alkyl, C_{1-3} alkylphenyl, wherein the phenyl-group can be substituted with a
60 group OH, NO_2 , NH_2 , CN, CH_3 , Cl, Br;

61 R_5 , R_6 , R_7 , which may be the same or different from one another, represent a
62 hydrogen or C_{1-3} alkyl group; their pharmaceutically acceptable salts, their
63 enantiomers and mixture thereof.

1 2. Compounds according to Claim 1, in which:

2 f, g, h, m, which may be the same or different from one another, may be 0 or 1;

3 R_1 and R_2 , which may be the same or different from one another, represent the
4 side chain of a natural amino acid chosen from among tryptophan, phenyl
5 alanine, tyrosine, histidine or the side chain of a non-natural amino acid chosen
6 in the group:

7 tryptophan and phenyl alanine, either mono- or di-substituted with residues
8 chosen from among C_{1-3} alkyl or halo-alkyl, C_{1-3} alkoxy or amino-alkoxy,
9 halogen, OH, NH_2 , $NR_{13}R_{14}$, where R_{13} and R_{14} , which may be the same or
10 different from one another, represent a hydrogen or C_{1-3} alkyl group;

11 R_3 represents a group chosen from among:

12 - linear or branched alkyl having the formula C_nH_{2n+1} , with $n = 1-5$ (chosen in the
13 group consisting of methyl, ethyl, propyl, isopropyl, n-butyl, t-butyl) cycloalkyl or
14 alkylcycloalkyl of formula C_nH_{2n-1} with $n = 5-9$ (chosen in the group consisting of
15 cyclopentyl, cyclohexyl, methylcyclohexyl)

16 - $(CH_2)_r-Ar_1$, where $r = 1$ or 2 and where Ar_1 is an aromatic group chosen in the
17 group consisting of: α -naphthyl, β -naphthyl, phenyl, indole, the said Ar_1 group
18 being possibly substituted with a maximum of 2 residues chosen in the group
19 consisting of: C_{1-3} alkyl, CF_3 , C_{1-3} alkoxy, Cl, F, OH, NH_2 ;

20 R_4 represents an L-Q group where:

21 L is a chemical bond or CH_2 , and

22 Q is a group chosen from among:

23 - OH, NH_2 , NR_9R_{10} , OR_{11} , and where R_9 and R_{10} , which may be the same or
24 different from one another, represent a hydrogen or C_{1-3} alkyl group, C_{1-3} hydroxy
25 alkyl, C_{1-3} dihydroxyalkyl, C_{1-3} alkyl-CONHR₁₂ (wherein R_{12} is a monoglycosidic
26 group derived from D or L pentoses or hexoses (chosen in the group consisting
27 of ribose, arabinose, glucose, galactose, fructose, glucosamine, galactosamine
28 and their N-acetylated derivatives)), C_{1-3} alkyltetrazole, C_{1-3} alkyl-COOH or
29 wherein R_9R_{10} are joined together to form with the N atom a morpholine or a
30 piperidine ring and where R_{11} is a C_{1-3} alkyl chain, or a C_{2-4} amino-alkyl chain;

- 31 - NHCOR_8 wherein R_8 is a cyclohexane containing from 2 to 4 OH groups, a C_{1-6}
 32 alkylchain containing a polar group (chosen in the group consisting of NH_2 ,
 33 COOH , CONHR_{12} (wherein R_{12} is as hereabove define) or [1,4']bipiperidine)
 34 - COOH , COOR_{17} or CONHR_{12} , wherein R_{12} is as hereabove defined and R_{17} is
 35 as R_{12} or a group 4-nitrobenzyl.
 36 - R_5 , R_6 , R_7 are H.

37 in which the carbon atom that carries the substituents R_3 and R_7 has
 38 configuration R.

1 3. Compounds according to Claim 2, as specified below:

- 2 Cyclo{-Suc-Trp-Phe-[(R)-NH-CH($\text{CH}_2\text{C}_6\text{H}_5$)- CH_2 -NH-]}
 3 Cyclo{-Suc-Trp-Phe-[(S)-NH-CH($\text{CH}_2\text{C}_6\text{H}_5$)- CH_2 -NH-]}
 4 Cyclo{-Suc-Trp-Phe-[(R)-NH-CH($\text{CH}_2\text{C}_6\text{H}_{11}$)- CH_2 -NH-]}
 5 Cyclo{-Suc-Trp-Phe-[(R)-NH-CH($\text{CH}_2\text{C}_6\text{H}_4(4\text{-OCH}_3)$)- CH_2 -NH-]}
 6 Cyclo{-Suc-Trp(5F)-Phe-[(R)-NH-CH($\text{CH}_2\text{C}_6\text{H}_5$)- CH_2 -NH-]}
 7 Cyclo{-Suc-Trp(Me)-Phe-[(R)-NH-CH($\text{CH}_2\text{C}_6\text{H}_5$)- CH_2 -NH-]}
 8 Cyclo{-Suc-Phe(3,4-Cl)-Phe-[(R)-NH-CH($\text{CH}_2\text{C}_6\text{H}_5$)- CH_2 -NH-]}
 9 Cyclo{-Suc-Trp-Phe(3,4-Cl)-[(R)-NH-CH($\text{CH}_2\text{C}_6\text{H}_5$)- CH_2 -NH-]}
 10 Cyclo{-Suc-Trp-Tyr-[(R)-NH-CH($\text{CH}_2\text{C}_6\text{H}_5$)- CH_2 -NH-]}
 11 Cyclo{-Suc-Trp-Phe-[(R)-NH-CH($\text{CH}_2\text{C}_6\text{H}_3$ -3,4-diCl)- CH_2 -NH-]}
 12 Cyclo{-Suc-Trp-Phe-[(R)-NH-CH($\text{CH}_2\text{C}_6\text{H}_4$ -4-OH)- CH_2 -NH-]}
 13 Cyclo{-Suc-Trp-Phe-[(R)-NH-CH($\text{CH}_2\text{-CH}_2\text{-C}_6\text{H}_5$)- CH_2 -NH-]}
 14 Cyclo{-Suc-Trp-Phe-[(R)-NH-CH(CH_2 -2-naphthyl)- CH_2 -NH-]}
 15 Cyclo{-Suc-Trp-Phe-[(R)-NH-CH(CH_2 -indol-3-yl)- CH_2 -NH-]}
 16 Cyclo{-Suc-Trp-Phe-[(R)-NH-CH(CH_2 -5-F-indol-3-yl)- CH_2 -NH-]}
 17 Cyclo{-Suc-Trp-Phe-[(R)-NH-CH($\text{CH}_2\text{C}_6\text{H}_4$ -3-F)- CH_2 -NH-]}
 18 Cyclo{-Suc-Trp-Phe-[(R)NH-CH($\text{CH}_2\text{-C}_6\text{H}_3$ -3,4-diF- CH_2 -NH-)-]}
 19 Cyclo{-Suc-Trp-Phe-[(R)NH-CH($\text{CH}_2\text{-C}_6\text{H}_4$ -4- CF_3 - CH_2 -NH-)-]}
 20 Cyclo{-Suc-Trp-Phe-[(R)-NH- CH_2 -CH($\text{CH}_2\text{C}_6\text{H}_5$)-NH-]}
 21 Cyclo{-Suc-Trp-Phe-[(S)-NH- CH_2 -CH($\text{CH}_2\text{C}_6\text{H}_5$)-NH-]}
 22 Cyclo{-Trp-Phe-[(R)-NH-CH($\text{CH}_2\text{-C}_6\text{H}_5$)- CH_2 -NH-]}-(CH_2)₃CO-}
 23 Cyclo{-Trp-Phe-[(R)-NH-CH($\text{CH}_2\text{-C}_6\text{H}_5$)- CH_2 -N(CH_3)]-(CH_2)₃CO-}
 24 Cyclo{-Suc[1/(S)-NH]-Trp-Phe-[(R)-NH-CH($\text{CH}_2\text{C}_6\text{H}_5$)- CH_2 -NH-]}-(CH_2)₃CO-}


- 25 Cyclo{-Suc[1(R)-NH₂]-Trp-Phe-[(R)NH-CH(CH₂-C₆H₅)-CH₂NH]-}
- 26 Cyclo{-Suc[2(S)-NH₂]-Trp-Phe-[(R)NH-CH(CH₂-C₆H₅)-CH₂NH]-}
- 27 Cyclo{-Suc[2(R)-NH₂]-Trp-Phe-[(R)NH-CH(CH₂-C₆H₅)-CH₂NH]-}
- 28 Cyclo{-Suc[1(S)-NH(CH₃)]-Trp-Phe-[(R)NH-CH(CH₂-C₆H₅)-CH₂NH]-}
- 29 Cyclo{-Suc[1-COO(CH₂-C₆H₄-4-NO₂)]-Trp-Phe-[(R)NH-CH(CH₂-C₆H₅)-CH₂NH]-}
- 30 Cyclo{-Suc(1-COOH)-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂-NH]-}
- 31 Cyclo{-Suc(1-COOH)-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂-NH]-}
- 32 Cyclo{-Suc(1-OH)-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂-NH]-}
- 33 Cyclo{-Suc(2-COOH)-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂-NH]-}
- 34 Cyclo{-Suc(2-OH)-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂-NH]-}
- 35 Cyclo{-Suc[1(S)-(2H-tetrazolyl-5-ylmethyl)amino]-Trp-Phe-[(R)-NH-CH(CH₂-
- 36 C₆H₅)-CH₂-NH]-}.TFA
- 37 Cyclo{-Suc[1(S)-(morpholin-4-yl)]-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂-NH]-
- 38 }.TFA
- 39 Cyclo{-Suc[1(S)-N(CH₃)₂]-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂NH]-}.TFA
- 40 Cyclo{-Suc[1(S)-(piperidin-4-yl)]-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂NH]-}.TFA
- 41 Cyclo{-Suc[1(S)-(N(CH₂CH₂OH)₂)]-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂-
- 42 NH]-}.TFA
- 43 Cyclo{-Suc[1(S)-(N(CH₂CH(OH)CH₂OH)]-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂-
- 44 NH]-}.TFA
- 45 Cyclo{-Suc[1(S)-(3-carboxypropanoyl)amino]-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-
- 46 CH₂-NH]-}.
- 47 Cyclo{-Suc[1(S)-[3-N'-(β-D-glucopiranos-1-yl)-carboxamidopropanoyl]amino]-
- 48 Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂-NH]-}
- 49 Cyclo{-Suc[1(S)-[(carboxymethyl)amino]-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂-
- 50 NH]-} TFA
- 51 Cyclo{-Suc[1(S)-[N'-(β-D-glucopiranos-1-yl)-carboxyamidomethyl]amino]-Trp-
- 52 Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂-NH]-} TFA
- 53 Cyclo{-Suc[1(S)-(chinyll)amine]-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂-NH]-}
- 54 Cyclo{-Suc[1(S)-(4-aminobutanoyl)amino]-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂-
- 55 NH]-} TFA

- 56 Cyclo{-Suc[1(S)-[(1,4')bipiperidin-1-yl]acetamido]-Trp-Phe-[(R)-NH-CH(CH₂-
 57 C₆H₅)-CH₂-NH]-} TFA
 58 Cyclo{-Suc[1-N-(β-D-glucopiranos-1-yl)-carboxyamido]-Trp-Phe-[(R)-NH-
 59 CH(CH₂-C₆H₅)-CH₂-NH]-}
 60 Cyclo{-Suc[1(S)-[N'-(2-N-acetyl-β-D-glucopiranos-1-yl)-carboxyamido]-Trp-Phe-
 61 [(R)-NH-CH(CH₂-C₆H₅)-CH₂-NH]-}.


- 1 4. Process for the synthesis of a compound of general formula (I), where X₁, X₂,
 2 X₃, X₄ are CONH and the other substituents are as defined in Claim 1, where:
 3 a) the suitably protected amino acids (1), (2) and (4)



(1)




(2)




(4)

- 9 are made to react, as shown in the diagram, with the derivative of the protected
 10 succinic acid (7)




(7)

- 16 thus obtaining the linear compound (8)



(8)

- 21 b) the linear compound 8, is deprotected and cyclicized to yield the final
 22 monocyclic compound (10)



- 1 5. Pharmaceutical compositions containing as active principle the compounds
2 of general formula (I) according to Claim 1 in combination with pharmaceutically
3 acceptable carriers or excipients.
- 1 6. Pharmaceutical compositions according to Claim 5, to be used as tachykinin
2 antagonists.
- 1 7. Pharmaceutical compositions according to Claim 6, to be used as
2 antagonists of the human NK-2 receptor.
- 1 8. Pharmaceutical compositions according to Claim 7, to be used in the
2 treatment of the bronchospastic and inflammatory component of asthma,
3 coughing, pulmonary irritation, intestinal spasms, spasms of the biliary tract,
4 local spasms of the bladder and of the ureter during cystitis, and kidney
5 infections and colics.
- 1 9. Pharmaceutical compositions according to Claim 7, to be used as anxiolytics.
- 1 10. Use of a compound according to Claim 1 as tachykinin antagonist.
- 1 11. Use of a compound according to Claim 1 as NK-2 antagonist.
- 1 12. Use of a compound according to Claim 1 in the treatment of the
2 bronchospastic and inflammatory component of asthma, coughing, pulmonary
3 irritation, intestinal spasms, spasms of the biliary tract, local spasms of the
4 bladder and of the ureter during cystitis, and kidney infections and colics.
- 1 13. Use of a composition according to Claim 1 as an NK-2 antagonist for the
2 treatment of anxiety syndromes.

1 14. Method for the treatment of the bronchospastic and inflammatory
2 component of asthma, coughing, pulmonary irritation, intestinal spasms,
3 spasms of the biliary tract, local spasms of the bladder and of the ureter during
4 cystitis, and kidney infections and colics, in which quantities of between 0.02
5 and 10 mg/kg of body weight of active principle consisting of products of
6 formula (I), according to Claim 1, are administered to the patient.



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07K 5/065, 7/54	A3	(11) International Publication Number: WO 98/34949 (43) International Publication Date: 13 August 1998 (13.08.98)
(21) International Application Number: PCT/EP98/00599 (22) International Filing Date: 4 February 1998 (04.02.98) (30) Priority Data: FI97A000020 7 February 1997 (07.02.97) IT (71) Applicant (for all designated States except US): MENARINI RICERCHE S.p.A. [IT/IT]; Via Tito Speri, 10, I-00040 Pomezia (IT). (72) Inventors; and (75) Inventors/Applicants (for US only): GIORGI, Raffaello [IT/IT]; Via Delle Piagge, 9, I-56124 Pisa (IT). DI BUGNO, Cristina [IT/IT]; Via R. Sanzio, 16, I-56122 Pisa (IT). GIANNOTTI, Danilo [IT/IT]; Via Roma, 128, I-55011 Altopascio (IT). MAGGI, Carlo, Alberto [IT/IT]; Via Michelazzi, 43, I-50141 Firenze (IT). (74) Agent: PASSINI, Angelo; Notarbartolo & Gervasi S.p.A., Corso di Porta Vittoria, 9, I-20122 Milan (IT).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> (88) Date of publication of the international search report: 22 October 1998 (22.10.98)
(54) Title: MONOCYCLIC COMPOUNDS WITH FOUR BIFUNCTIONAL RESIDUES HAVING NK-2 ANTAGONIST ACTION (57) Abstract <p>The present invention refers to compounds of general formula (I) having NK-2 antagonist action, pharmaceutical compositions containing them, and processes for their preparation.</p> <div style="text-align: center;"> <p style="text-align: right;">(I)</p> </div>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

INTERNATIONAL SEARCH REPORT

National Application No

PCT/EP 98/00599

A. CLASSIFICATION OF SUBJECT MATTER
 IPC 6 C07K5/065 C07K7/54

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
 IPC 6 C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 4 703 034 A (FREIDINGER ROGER ET AL) 27 October 1987 see column 11; claim 1; table IV	1,2
X	KITABATAKE K. ET AL.: "GUSHING- INDUCING PEPTIDES IN BEER PRODUCED BY PENICILLUM CHYRSOGENUM" PEPT.CHEM, vol. 17, 1980, TOKYO, pages 7-12, XP002073620 see table 3	1,2
Y	WO 96 28467 A (MENARINI FARMA IND ; ARCAMONE FEDERICO (IT); MAGGI CARLO ALBERTO (I) 19 September 1996 see claim 1	1-14
	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance
 "E" earlier document but published on or after the international filing date
 "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
 "O" document referring to an oral disclosure, use, exhibition or other means
 "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
 "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
 "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
 "&" document member of the same patent family

Date of the actual completion of the international search

4 August 1998

Date of mailing of the international search report

14/08/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Deffner, C-A

INTERNATIONAL SEARCH REPORT

Inventor's Application No

PCT/EP 98/00599

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EP 0 333 174 A (FUJISAWA PHARMACEUTICAL CO) 20 September 1989 see claim 1 -----	1-14

INTERNATIONAL SEARCH REPORT

International Application No

Information on patent family members

PCT/EP 98/00599

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 4703034	A	27-10-1987	NONE	
<hr/>				
WO 9628467	A	19-09-1996	IT FI950044 A	13-09-1996
			AU 5105996 A	02-10-1996
			BR 9607348 A	30-12-1997
			CA 2215372 A	19-09-1996
			CZ 9702862 A	18-02-1998
			EP 0815126 A	07-01-1998
			HR 960117 A	31-08-1997
			NO 974057 A	07-11-1997
			PL 322105 A	05-01-1998
			SK 121297 A	04-02-1998
<hr/>				
EP 0333174	A	20-09-1989	AT 137763 T	15-05-1996
			AU 3132489 A	21-09-1989
			CA 1329444 A	10-05-1994
			CN 1037156 A	15-11-1989
			DE 68926403 D	13-06-1996
			DE 68926403 T	17-10-1996
			DK 126389 A	17-09-1989
			FI 891176 A	17-09-1989
			JP 1227025 A	17-11-1988

